

FEDAC

PRODUCT DESCRIPTION:

Tablets : Light pink, round and flat tablets, 8 mm diameter with score line and "DHA" logo.
Syrup : Clear mixture, yellow in colour, sweet taste with orange flavour.

COMPOSITION:

1) Tablets: Tripolidine HCl BP 2.5mg/tablet
Pseudoephedrine HCl BP 60mg/tablet
2) Syrup : Tripolidine HCl BP 1.25mg/5ml
Pseudoephedrine HCl BP 30mg/5ml

PRESERVATIVES:

Methyl hydroxybenzoate 0.15% w/v
Propyl hydroxybenzoate 0.015% w/v

PHARMACOLOGY:

Tripolidine is an antihistamine which competes with histamine for H₁-receptor sites on effector cells. It prevents but does not reverse responses mediated by histamine alone. Its anti-cholinergic actions provide a drying effect on the nasal mucosa. Pseudoephedrine acts on alpha-adrenergic receptors in the mucosa of the respiratory tract, producing vasoconstriction. It shrinks swollen nasal mucous membrane, reduces tissue hyperemia, oedema and nasal congestion, and increases nasal airway patency. Also drainage of sinus secretions may be increased and obstructed eustachian ostia may be opened.

INDICATIONS:

Symptomatic treatment of nasal and respiratory congestion, common cold, acute sinusitis, allergic rhinitis.

RECOMMENDED DOSAGE:

Tablet : Adults : 1 tablet 3 times a day

Syrup : Adults and children over 12 years: 2 teaspoonfuls 3 times a day.

Children : 6 - 12 years: 1 teaspoonful,
2 - 5 years: 1/2 teaspoon.

Doses to be taken 3 times a day.

Note: 1 teaspoonful is equal to 5ml.

CAUTION:

Not to be used in children less than 2 years of age.

To be used with caution and on doctor's or pharmacist's advice in children 2 to 6 years of age.

ADVERSE EFFECTS:

Drowsiness, dryness of mouth, nervousness, restlessness, trouble in sleeping.

PRECAUTIONS:

Patients sensitive to one antihistamine or sympathomimetic may be sensitive to others. Tripolidine may cause drowsiness. If affected do not drive or operate machinery.

USE IN PREGNANCY:

There are no specific data on the use of tripolidine and pseudoephedrine during pregnancy. Caution should therefore be exercised by weighing the potential benefits against the possible risks to the foetus before prescribing the drug.

CONTRAINDICATIONS:

Individuals who are hypersensitive to any component in this product, patients with severe hypertension or severe coronary artery disease; patients who are taking or have taken MAOIs within the preceding 2 weeks; patients with hyperthyroidism.

DRUG INTERACTIONS:

Concomitant use of Fedac with sympathomimetic agents or with MAOIs may cause a rise in blood pressure. Pseudoephedrine may reverse the hypotensive activity of drugs which interfere with sympathetic activity. Tripolidine may enhance the drowsiness caused by sedative drugs and alcohol.

SYMPTOMS AND TREATMENT OF OVERDOSAGE:

Symptoms: Restlessness, delirium and muscle tremors.

Treatment: Give activated charcoal and then remove the drug by airway protected gastric lavage followed by catharsis.

AVAILABILITY:

Tablets : 10 x 10 x 10's blister pack

Syrup : 3.8 litres in high density polyethylene container with cap.
120ml in amber glass bottle.

Not all pack sizes may be available locally.

EXPIRY PERIOD AND STORAGE:

Do not use after the expiry date stated on the label. Store below 30°C and protect from light.

DATE OF REVISION: September 2020

For further information, please consult your physician or pharmacist.

Manufactured by:
PT Actavis Indonesia
Jl. Raya Bogor Km 28
Jakarta
Indonesia, 13710

41095-03

		Product Name		Leaflet Fedac Tablet and Syrup		Scale : 100%		
		Dimension		120 x 208 mm		Color White block		
		Item Code		41095-03				
(I) Supplier must send the Proof Printing material within 14 days after Actavis Indonesia send the AW final approval (I)		<u>Specification :</u> - Material : HVS 70gsm - Type of Varnish (for Carton) : - Lipat / Not (for Dirslip) : - Etc : 				Market : Singapore		
R E V I S I O N	Version	Change Classified	Description of change			Date	Issued by	Approved by
	02 02		remove product owner’s address n chinese characters change storage to 30			July 7, 20 Sept 24, 20	market market	
ACTAVIS						MARKET APPROVAL		
REG		PM	PROD	QA	QC			
Date :		Date :	Date :	Date :	Date :			
Initials :		Initials :	Initials :	Initials :	Initials :			